

# Endarterectomy for Asymptomatic Carotid Artery Stenosis

Executive Committee for the Asymptomatic Carotid Atherosclerosis Study

**Objective.**—To determine whether the addition of carotid endarterectomy to aggressive medical management can reduce the incidence of cerebral infarction in patients with asymptomatic carotid artery stenosis.

**Design.**—Prospective, randomized, multicenter trial.

**Setting.**—Thirty-nine clinical sites across the United States and Canada.

**Patients.**—Between December 1987 and December 1993, a total of 1662 patients with asymptomatic carotid artery stenosis of 60% or greater reduction in diameter were randomized; follow-up data are available on 1659. At baseline, recognized risk factors for stroke were similar between the two treatment groups.

**Intervention.**—Daily aspirin administration and medical risk factor management for all patients; carotid endarterectomy for patients randomized to receive surgery.

**Main Outcome Measures.**—Initially, transient ischemic attack or cerebral infarction occurring in the distribution of the study artery and any transient ischemic attack, stroke, or death occurring in the perioperative period. In March 1993, the primary outcome measures were changed to cerebral infarction occurring in the distribution of the study artery or any stroke or death occurring in the perioperative period.

**Results.**—After a median follow-up of 2.7 years, with 4657 patient-years of observation, the aggregate risk over 5 years for ipsilateral stroke and any perioperative stroke or death was estimated to be 5.1% for surgical patients and 11.0% for patients treated medically (aggregate risk reduction of 53% [95% confidence interval, 22% to 72%]).

**Conclusion.**—Patients with asymptomatic carotid artery stenosis of 60% or greater reduction in diameter and whose general health makes them good candidates for elective surgery will have a reduced 5-year risk of ipsilateral stroke if carotid endarterectomy performed with less than 3% perioperative morbidity and mortality is added to aggressive management of modifiable risk factors.

(*JAMA*. 1995;273:1421-1428)

tion in individuals with asymptomatic hemodynamically significant carotid artery stenosis?

For editorial comment see p 1459.

Secondary objectives were to determine the surgical success in lesion removal and the incidence of recurrent carotid stenosis, the rate of progression or regression of carotid atherosclerosis in the medically treated comparison group, and the incidence of all other vascular events, such as TIA, myocardial infarction, and death related to vascular disease during follow-up.

## METHODS

The design and organization of the ACAS are detailed elsewhere.<sup>13</sup> Thirty-nine clinical centers were chosen from an applicant pool of 55. All obtained institutional review board approval of the study protocol.

## Recruitment

Study participants were recruited from ultrasound vascular laboratories, practitioners who auscultated carotid bruits, and physicians who found carotid stenosis during evaluation for peripheral vascular surgery or contralateral CEA.

Inclusion criteria were age between 40 and 79 years; compatible history and findings on physical and neurological examinations; performance of required laboratory and electrocardiographic examinations no earlier than 3 months before randomization; patient accessibility and willingness to be followed for 5 years; and valid informed consent.

Exclusion criteria were cerebrovascular events in the distribution of the study carotid artery or in that of the vertebrobasilar arterial system; symptoms referable to the contralateral cerebral hemisphere within the previous 45 days; contraindication to aspirin therapy; a disorder that could seriously complicate surgery; or a condition that could prevent continuing participation or was likely to produce disability or death within 5 years. (Detailed information regarding eligibil-

MORE THAN 500 000 new strokes occur annually in the United States, and it has been estimated that carotid artery disease may be responsible for 20% to 30% of them.<sup>1</sup> The annual stroke event rate for asymptomatic patients with hemodynamically significant carotid artery stenosis ranges from 2% to 5%.<sup>2-5</sup> Carotid artery stenosis usually is identified after transient ischemic attack (TIA), but for many patients, cerebral infarction caused by artery-to-artery embolism or carotid occlusion is the initial event. Progression of asymptomatic carotid artery stenosis to occlusion is unpredictable and can be disastrous; at the time of occlusion, dis-

abling stroke may occur in 20% of patients, and thereafter in 1.5% to 5% annually.<sup>6-8</sup> On the other hand, the 30-day major morbidity and mortality for patients who undergo surgery for asymptomatic stenosis ranges from 0.0% to 3.8%, and that for patients with symptomatic stenosis may be 6%.<sup>9</sup> Because the role of carotid endarterectomy (CEA) for asymptomatic carotid artery stenosis had not been proved,<sup>10-12</sup> the Asymptomatic Carotid Atherosclerosis Study (ACAS) was initiated in 1987.<sup>13</sup> The ACAS is an investigator-initiated randomized trial designed to test whether CEA should be a component of management for selected patients with asymptomatic stenosis of the common carotid bulb, the internal carotid sinus, or both. The question addressed was: Will CEA added to aggressive reduction of modifiable risk factors and administration of aspirin reduce the 5-year risk of ipsilateral cerebral infarc-

A complete list of the collaborators in the Asymptomatic Carotid Atherosclerosis Study appears at the end of this article.

Reprint requests to Stroke Center and Department of Neurology, Bowman Gray School of Medicine of Wake Forest University, Medical Center Blvd, Winston-Salem, NC 27157-1068 (James F. Toole, MD).

L



eds  
on't  
cation.

the same.  
erapies and

TYLENOL



d risks.

ity and exclusion is available on request from the corresponding author.)

The ACAS definition of hemodynamically significant carotid stenosis required that at least one of three criteria was met: arteriography within the previous 60 days indicating stenosis of at least 60% reduction in diameter (if the arteriogram was performed 61 to 364 days before randomization, Doppler ultrasonography was required to verify that the artery had not occluded); Doppler examination within the preceding 60 days showing a frequency or velocity greater than the instrument-specific cut point with 95% positive predictive value (PPV); or Doppler examination showing a frequency or velocity greater than the instrument-specific 90% PPV cut point confirmed by ocular pneumoplethysmographic (OPG-Gee) examination performed within the previous 60 days.

A patient could enter the study with unilaterally or bilaterally asymptomatic, hemodynamically significant stenosis, but only one artery was the study artery. If two arteries were eligible, the one with the greater stenosis was selected. If the stenoses were identical, the left carotid artery was chosen. Patients randomized to surgery on the basis of Doppler or Doppler with OPG-Gee were required to have an arteriogram prior to CEA. If a postrandomization arteriogram revealed the contralateral artery to have the greater stenosis, it then became the study artery. The nonstudy artery was managed medically unless a cerebrovascular event occurred, at which time CEA could be considered.

### Arteriographic Measurements

The minimal residual lumen (MRL) and the distal lumen (DL) were measured on the same radiograph. The MRL was the smallest lumen diameter at the site of the stenotic lesion. The DL was the diameter at the first point distal to the MRL at which the arterial walls became parallel. Percentage of stenosis was calculated as  $100 \times (1 - [MRL/DL])$ .

### Ultrasound Measurements

Because of the heterogeneity among ultrasound devices and techniques, we established a cut point for each by comparing Doppler ultrasonography with arteriograms performed within 42 days of each other. Doppler cut points were computed for peak systolic frequency or, if indeterminate, end diastolic frequency, based on data from 50 consecutive patients.<sup>14</sup>

### Randomization

An ACAS neurologist and an ACAS surgeon gave joint approval for entering patients. Once the eligibility criteria had been confirmed and after informed consent was obtained, the patient was

randomized using the permuted block method with at least three different block sizes determined randomly, stratified by center, gender, number of eligible arteries, and previous contralateral CEA. The assignment category was communicated to each clinical center by the statistical coordinating center through an individualized computer program arranged so that the clinical center could neither predict nor reject an assignment.

### Medical Treatment

All patients received 325 mg of regular or enteric-coated aspirin daily (provided by Sterling Health USA, New York, NY). Stroke risk factors and their modification were reviewed with all patients at the time of randomization and again during subsequent interviews and telephone follow-up. This included discussion of diastolic and systolic hypertension, diabetes mellitus, abnormal lipid levels, excessive consumption of ethanol, and tobacco use. Whenever possible, the recommendations of the ACAS Risk Factor Reduction Committee were followed (available on request from the corresponding author).

### Surgical Treatment

In addition to 325 mg of aspirin daily and risk factor modification counseling, patients randomized to the surgical arm received the normal evaluation and care of a surgical patient. They were scheduled to undergo CEA within 2 weeks of randomization. If an arteriogram or cranial computed tomogram (CCT) had not been performed, the patient underwent the procedure(s) before CEA. The arteriogram must have demonstrated a stenosis of 60% or greater. Patients with a postrandomization, presurgery arteriogram demonstrating less than 60% stenosis or a distal abnormality such as aneurysm, arteriovenous malformation, or siphon stenosis exceeding the proximal stenosis did not undergo surgery but were retained in the surgical arm for comparison analyses. Asymptomatic cerebral infarction demonstrated by CCT was not an exclusion for surgery.<sup>15</sup> No attempt was made to standardize or control anesthesia or surgical techniques used by the 117 ACAS-credentialed surgeons.<sup>16</sup>

The surgeon, the ACAS neurologist, and the ACAS patient coordinator examined each patient 24 hours after CEA. All deficits occurring through the 30-day perioperative period required the administration of the end point review process (described below).

### Follow-up

Follow-up evaluations were conducted at 1 month and thereafter every 3 months, alternating between clinic visits and telephone contacts. During the clinic visit,

patients completed a medical history questionnaire and TIA/stroke questionnaire and underwent physical and neurological examinations and a Mini-Mental State Examination.<sup>17</sup> Risk reduction management was reviewed and aspirin adherence was determined by pill count.

Doppler ultrasound studies were repeated at the 3-month follow-up, every 6 months thereafter during the first 24 months, then yearly, and at potential or verified end point or at exit from the study after 5 years; CCT was repeated at potential end point or exit. Electrocardiogram was repeated when clinically indicated and at exit.

Patients were instructed to notify the coordinator if symptoms suggesting possible TIA or stroke occurred. The coordinator scheduled urgent evaluations by the ACAS neurologist and surgeon, and activated the end point verification system.

In addition to identification of events from clinic visits and telephone contacts, hospital discharge diagnoses and death certificates were reviewed for coronary events and strokes.

### End Point Definition and Verification

A TIA was defined as a focal ischemic neurological deficit of abrupt onset lasting at least 30 seconds and resolving completely within 24 hours. Deficits persisting longer than 24 hours were classified as stroke.<sup>18</sup> All strokes or deaths occurring within 30 days after randomization in the surgical and 42 days in the medical groups were included as end points to reflect operative morbidity and mortality. The difference in times reflected an average 12-day interval between randomization and surgery.

Initial review was conducted under a stringent timetable involving one external expert masked for local diagnosis, treatment assignment, clinical center, and temporal relationship to surgery (if done). In addition, every potential event was abstracted, masked, and reviewed together by all six experts on the End Point Review Committee. Our analyses are based on their diagnoses.

Secondary analyses considered any stroke and perioperative death; any stroke and any death; and any ipsilateral TIA and stroke and any perioperative TIA, stroke, or death. The ACAS used categories 2 through 5 of the Glasgow scale to determine a major stroke, defined as a stroke resulting in moderate or severe disability, persistent vegetative state, or death.<sup>19</sup>

### Statistical Analyses

Initially, the primary end points for evaluation of the two treatments were

Table 1.—Baseline Characteristics of Patients by Treatment

Baseline Characteristic	
Age, y	
40-49	
50-59	
60-69	
70-79	
Race	
White	
Black	
Other	
Sex	
M	
F	
History	
Coronary artery disease	
Hypertension†	
Cancer	
Diabetes mellitus	
Lung disease at entry	
Current cigarette smoker	
Bilaterally eligible artery	
Previous contralateral CEA	
Endarterectomy	
TIA or stroke§	
EC-IC bypass	
Subclavian bypass	
Bruits in neck	
Ipsilateral carotid stenosis	
Contralateral carotid stenosis	
Infarct on CT scan¶	
Any location	
Ipsilateral silent infarct	
Contralateral occlusion by Doppler	

\*TIA indicates transient ischemic attack.  
 †TIA indicates transient ischemic attack.  
 ‡Defined as positive arteriogram, previous myocardial infarction, or previous stroke.  
 §Positive response to treatment that you had high blood pressure.  
 ¶Significant difference in blood pressure (unadjusted for multiple comparisons).  
 ||Bruit question not asked of medical patients. Ipsilateral carotid stenosis.  
 ¶CT scan was not performed on medical patients.

ipsilateral TIA and stroke. In a multivariate analysis, the risk of cerebrovascular events was similar in both groups. Because of the wide use, a two-sided hypothesis ( $\alpha = 0.05$ ) calculations including ipsilateral TIA and stroke according to our definitions (intentionally assuming an annual 10% outcome group for cerebral infarction) indicated that 75% of each treatment group could detect a 35% difference in rates, allowing for a 10% follow-up.

The results of this trial<sup>20</sup> demonstrate that preventing TIA and carotid stenosis, and that automatic Carotid Endarterectomy demonstrated that TIA is better

Table 1.—Baseline Characteristics of Randomized Patients by Treatment Assignment, in Percentages\*

Baseline Characteristic	Treatment Assignment	
	Surgical (n=825)	Medical (n=834)
Age, y		
40-49	2	2
50-59	13	15
60-69	50	46
70-79	36	38
Race		
White	94	95
Black	3	3
Other	3	2
Sex		
M	66	66
F	34	34
History		
Coronary artery disease†	69	69
Hypertension‡	64	64
Cancer	12	10
Diabetes mellitus	25	21
Lung disease at entry	6	5
Current cigarette smoker	28	24
Bilaterally eligible arteries	10	9
Previous contralateral Endarterectomy	20	19
TIA or stroke§	22	27
EC-IC bypass	<1	0
Subclavian bypass	<1	<1
Bruits in neck¶		
Ipsilateral	76	74
Contralateral	44	42
Infarct on CT scan¶		
Any location	22	24
Ipsilateral silent	8	9
Contralateral occlusion by Doppler	10	9

\*TIA indicates transient ischemic attack; EC-IC, external carotid-internal carotid; and CT, computed tomographic.  
 †Defined as positive history of angina, coronary artery bypass, previous myocardial infarction, or abnormal electrocardiogram.  
 ‡Positive response to "Has your doctor ever told you that you had high blood pressure or were hypertensive?"  
 §Significant difference between groups at .05 level (unadjusted for multiple comparisons).  
 ¶Bruit question not evaluated for 51 surgical and 52 medical patients. Ipsilateral/contralateral refers to randomized artery.  
 ¶CT scan was not available on 71 surgical and 60 medical patients.

ipsilateral TIA, stroke, or any perioperative TIA, stroke, or death. The primary analysis is the comparison of the 5-year risk of cerebrovascular events in the two groups. Because both treatments were in wide use, a two-sided test of the null hypothesis ( $\alpha=.05$ ) was chosen. Power calculations included all randomized patients according to original treatment assignments (intention-to-treat analysis). Assuming an annual event rate in the worse-outcome group of 3% for TIA and 1% for cerebral infarction, these calculations indicated that 750 patients were needed in each treatment arm for 90% power to detect a 35% difference in 5-year event rates, allowing for as much as 20% loss to follow-up.

The results of the Veterans Affairs trial<sup>20</sup> demonstrated that CEA is preferable to medical management for preventing TIA in asymptomatic carotid stenosis, and the North American Symptomatic Carotid Endarterectomy Trial<sup>21</sup> demonstrated that infarction following TIA is better managed surgically. Nei-

ther resolved the issue of whether CEA prevents unheralded cerebral infarction. Therefore, in March 1993, the ACAS Executive Committee and the Data and Safety Monitoring Committee voted to restrict the primary end point to stroke and perioperative complications or death. For baseline comparisons, we used two-tailed *t* tests for comparing the means of continuous variables and  $\chi^2$  for comparing distributions of categorical variables, with no adjustment for multiple comparisons. Kaplan-Meier estimates of 5-year aggregate risk were compared between treatment groups using either Greenwood's formula for variances, for a large-sample test ignoring randomization stratification,<sup>22</sup> or randomization tests, respecting randomization strata.<sup>23</sup> In the initial years of treatment comparison, 1991 through 1993, the tests were for 2-, 3-, or 4-year aggregate risk. The randomization test was the primary method for interim treatment comparisons (see below). By the time of study closure *P* values from the two methods agreed within .002, so that all test results and confidence intervals (CIs) reported are based on large-sample tests unless otherwise noted.

Semiannual treatment comparison analyses were used to advise the Data Safety and Monitoring Committee whether a significance boundary had been crossed. The stopping rule was a modified O'Brien-Fleming<sup>24</sup> rule for maintaining the desired overall significance level despite repeated testing. The modification was for testing at selected intervals rather than predetermined numbers of events. The five analyses originally planned were changed to 10, because recruitment lagged and TIA was deleted as a primary end point. The critical value for the test statistic was 6.00 at the first test and 2.07 for the 10th, as compared with 1.96 for a single test at the significance level .05. The study was stopped after the eighth test, when the critical value was 2.38, corresponding to a nominal significance level of .017.

Treatment comparisons are reported herein in terms of relative risk reduction, the 5-year risk reduction due to surgery as a proportion of risk in the medical group.<sup>25</sup> Absolute treatment

Table 2.—Arteriographic Stenosis of the Ipsilateral Carotid Artery

% Stenosis	Prerandomization Arteriogram			Postrandomization Presurgery Arteriogram
	Medical	Surgical	Total (%)	
0-59	Not applicable	NA	NA	32 (8)
60-69	131	137	268 (42)	139 (34)
70-79	94	93	187 (29)	110 (26)
80-89	75	79	154 (24)	107 (26)
90-99	13	20	33 (5)	24 (6)
Total	313	329	642*	412†

\*Two patients were missing, one in each group.  
 †Two patients were missing.

group-specific risk levels are also provided for calculation of absolute risk reduction. For the primary event the number of patients treated to prevent one event over 5 years was calculated as the inverse of the absolute risk reduction.<sup>25</sup> Intention-to-treat analyses were used for all comparisons unless otherwise indicated, regardless of postrandomization ineligibility or crossover. End points for medical patients who received CEA after verified ipsilateral end point were categorized as perioperative if they occurred within 30 days of CEA. All tests were two tailed.

**COHORT CHARACTERISTICS**

During the 6 years of the study, more than 42 000 patients were screened and 1662 patients were randomized. Twenty-four centers contributed more than 30 patients each, and 13 contributed more than 50 patients. The average number of patients recruited per center was 43. From March 1988 through October 1993, 12 080 CEAs were performed at the sites. Six percent (683) were performed on "likely eligible nonrandomized" patients of ACAS physicians, 6% (758) were performed on already randomized ACAS patients, and the rest were performed on symptomatic patients, ineligible patients, or patients of surgeons not collaborating in the ACAS. Patient characteristics are presented in Table 1. Of the 1662 randomized patients, three in the surgery group were lost to follow-up after randomization and are excluded from analysis, leaving 1659. The 825 surgical and 834 medical patients were compared for 189 baseline characteristics, with only six tests yielding nominal statistically significant differences at the .05 level. Two thirds of the patients were men, 95% were white, and 48% were aged 60 through 69 years. Mean age was 67 years; mean weight, 81 kg for men and 67 kg for women; mean systolic blood pressure, 146 mm Hg; mean diastolic blood pressure, 78 mm Hg; and mean total cholesterol concentration, 5.90 mmol/L (228 mg/dL). Approximately 75% of patients had a bruit associated with the study artery, and in 43%, a contralateral carotid bruit was heard; 21% had a previous myo-

group-specific risk levels are also provided for calculation of absolute risk reduction. For the primary event the number of patients treated to prevent one event over 5 years was calculated as the inverse of the absolute risk reduction.<sup>25</sup> Intention-to-treat analyses were used for all comparisons unless otherwise indicated, regardless of postrandomization ineligibility or crossover. End points for medical patients who received CEA after verified ipsilateral end point were categorized as perioperative if they occurred within 30 days of CEA. All tests were two tailed.

group-specific risk levels are also provided for calculation of absolute risk reduction. For the primary event the number of patients treated to prevent one event over 5 years was calculated as the inverse of the absolute risk reduction.<sup>25</sup> Intention-to-treat analyses were used for all comparisons unless otherwise indicated, regardless of postrandomization ineligibility or crossover. End points for medical patients who received CEA after verified ipsilateral end point were categorized as perioperative if they occurred within 30 days of CEA. All tests were two tailed.

group-specific risk levels are also provided for calculation of absolute risk reduction. For the primary event the number of patients treated to prevent one event over 5 years was calculated as the inverse of the absolute risk reduction.<sup>25</sup> Intention-to-treat analyses were used for all comparisons unless otherwise indicated, regardless of postrandomization ineligibility or crossover. End points for medical patients who received CEA after verified ipsilateral end point were categorized as perioperative if they occurred within 30 days of CEA. All tests were two tailed.

group-specific risk levels are also provided for calculation of absolute risk reduction. For the primary event the number of patients treated to prevent one event over 5 years was calculated as the inverse of the absolute risk reduction.<sup>25</sup> Intention-to-treat analyses were used for all comparisons unless otherwise indicated, regardless of postrandomization ineligibility or crossover. End points for medical patients who received CEA after verified ipsilateral end point were categorized as perioperative if they occurred within 30 days of CEA. All tests were two tailed.

istory questionnaire neurological al State Examination difference was  
 s were re-up, every 6 he first 24 potential or t from the repeated at Electrocardiographically in-  
 o notify the resting position. The co-evaluations id surgeon, verification  
 n of events ie contacts, and death or coronary  
 al ischemic onset last-olving com-its persist-e classified aths occur-omization the medical d points to ind mortal-ected an en random-  
 ed under a one exter-diagnosis, center, and ry (if done). event was viewed to- End Point alyses are  
 dered any ; any stroke ateral TIA ative TIA, used cat-sgow scale efined as a or severe ve state, or  
 points for ents were

Table 3.—Number of Observed Events in Median 2.7-Year Follow-up, Estimated Number and Percentage of Events in 5 Years, Reduction Due to Surgery in 5-Year Risk as a Proportion of Risk in the Medical Group (95% CI), and Large-Sample *P* Value for Treatment Group Difference, by Event Type\*

Event Type	Medical (n=834)		Surgical (n=825)		Reduction Due to Surgery in 5-y Risk as a Proportion of Risk in Medical Group (95% CI)	<i>P</i>
	Observed No. of Events in Median 2.7-y Follow-up	Kaplan-Meier Estimate of 5-y Event Risk, No. (%)	Observed No. of Events in Median 2.7-y Follow-up	Kaplan-Meier Estimate of 5-y Event Risk, No. (%)		
Ipsilateral stroke or any perioperative stroke or death	52	92 (11.0)	33	42 (5.1)	0.53 (0.22 to 0.72)	.004
Major ipsilateral stroke or any perioperative major stroke or death	24	50 (6.0)	21	28 (3.4)	0.43 (-0.17 to 0.72)	.12
Ipsilateral TIA or stroke or any perioperative TIA or stroke or death	102	160 (19.2)	55	67 (8.2)	0.57 (0.39 to 0.70)	<.001
Any stroke or any perioperative death	86	146 (17.5)	60	102 (12.4)	0.29 (-0.05 to 0.52)	.09
Any major stroke or perioperative death	40	76 (9.1)	28	53 (6.4)	0.30 (-0.30 to 0.62)	.26
Any stroke or death	155	266 (31.9)	127	211 (25.6)	0.20 (-0.02 to 0.37)	.08
Any major stroke or death	116	213 (25.5)	100	171 (20.7)	0.19 (-0.08 to 0.39)	.16

\*CI indicates confidence interval; and TIA, transient ischemic attack.

cardial infarction, and 21% a previous coronary artery bypass. Sixty-four percent had hypertension, 26% were cigarette smokers, and 23% had diabetes mellitus.

Four hundred seven patients (25%) had had a previous hemispheric event contralateral to the study artery, and 1155 (70%) were asymptomatic in the distribution of both arteries.

Thirty-nine percent of patients were randomized on the basis of an arteriogram showing at least 60% stenosis of the carotid artery. Fifty-five percent were randomized with a Doppler PPV cut point of at least 95%, and 6% with a Doppler cut point of at least 90% confirmed by OPG-Gee. The positive predictive value of Doppler, estimated from the postrandomization presurgery angiogram, was 93%.

Table 2 shows the distribution of percentage of stenosis for prerandomization and postrandomization arteriograms before CEA. Because the health status of patients who received prerandomization arteriograms may differ from that of those who did not, a weighted estimate based on both categories is included. Five percent of patients had stenosis of the randomized artery less than 60%; 39%, 60% to 69% stenosis; 28%, 70% to 79% stenosis; 25%, 80% to 89% stenosis; and 5%, 90% to 99% stenosis, for a mean percentage of stenosis of 73%.

The Central Reading Center classified 536 baseline CCTs<sup>15</sup> as showing cerebral infarction. If an infarct was present, it was further classified by age, size, distribution, and volume. Local and central readers agreed on 89% of the cases. Using the Central Reading Center as the standard, the sensitivity of local reading was 71% and the specificity was 94%.

## RESULTS

Of the 825 surgical patients, 101 did not have ipsilateral arteriography or CEA, 45 because of patient refusal despite prior agreement to accept either treatment. Twelve patients were rejected for sur-

gery because of severe cardiac disease. Three had a stroke or died before arteriography or surgery was performed. Arteriograms found 33 patients to be ineligible, six because of intracranial abnormalities and 27 because of less than 60% carotid artery stenosis. Eight patients did not have surgery for various other reasons. Of the 834 patients randomized to medical treatment, 45 received CEA without a verified ipsilateral TIA or stroke. Thus, 146 (9%) patients did not receive the assigned treatment. Eleven patients dropped out from follow-up in the medical and nine in the surgical group.

During the perioperative period, 19 surgical patients (2.3%) had a stroke or died. Two patients had a stroke, one died prior to hospitalization, and five had a cerebral infarction as a direct result of arteriography, one of whom died. There were 10 nonfatal strokes and one fatal myocardial infarction during the 30-day postsurgery period. In the comparable perioperative period for the medical group, three patients (0.4%) had a cerebral infarction (two patients) or died (one patient). For the surgical group, the risk in the perioperative period was 2.3% (95% CI, 1.28% to 3.32%), whereas for the medical group it was 0.4% (95% CI, 0.0% to 0.8%).

All patients randomized to the surgical group were required to have arteriography. Of the 414 patients who underwent arteriography prior to CEA, five experienced a cerebral infarction, for an arteriographic complication rate of 1.2%. It is estimated that if all 724 patients receiving CEA had undergone arteriography as a part of the ACAS, 8.7 arteriographic cerebral infarctions would have occurred in addition to the 11 primary events in the 30 days following surgery, for an overall rate of 2.7% for cerebral infarction or death from the procedure.


Sixteen fatalities, potentially due to strokes, were reviewed by the Cerebrovascular End Point Review Committee. In no case was there a difference between the End Point Review Com-

mittee diagnosis and the local physician diagnosis. These events included six hemorrhagic strokes, two cerebral infarctions in the distribution of the randomized artery, three cerebral infarctions in the nonrandomized distribution, and five deaths not due to stroke.

## Treatment Comparisons

The study achieved its significance boundary after a median of 2.7 years of follow-up, with 9% of patients having completed 5 years; 26%, 4; 44%, 3; 68%, 2; and 87%, 1 year of follow-up. Because surgical patients were at greatest risk during the first month after endarterectomy, whereas the risk for medical patients was distributed throughout 5 years, comparisons near term greatly understated the differences expected after 5 years. Table 3 presents the observed number of events and also the Kaplan-Meier estimates predicted if all patients had been followed for 5 years. The estimated 5-year risk of ipsilateral stroke and any perioperative stroke or death was 11.0% for the medical group and 5.1% for the surgical group. The reduction in 5-year ipsilateral stroke risk in the surgical group was 53% of the estimated 5-year risk in the medical group (95% CI, 22% to 72%). The *P* value for the test of the difference between the treatment groups in 5-year risk of primary event was .004 by the large-sample test and the randomization test. For the primary end point of ipsilateral stroke and any perioperative stroke or death, the survival curves in the Figure cross near 10 months and become significantly reduced in the surgical group by 3 years (*P* < .05).

The results for secondary end points are in the same direction although not always statistically significant (Table 3 and Figure). Ipsilateral TIA or stroke or any perioperative TIA, stroke, or death—the original primary end point for the ACAS—showed a 57% reduction in 5-year risk for the surgery group (95% CI, 39% to 70%). In terms of any stroke or death,



Proportion of patients surviving without stroke or death from transient ischemic attack

the surgery group events (95% CI, 22% to 72%). The reduction in 5-year ipsilateral stroke risk in the surgical group was 53% of the estimated 5-year risk in the medical group (95% CI, 22% to 72%). The *P* value for the test of the difference between the treatment groups in 5-year risk of primary event was .004 by the large-sample test and the randomization test. For the primary end point of ipsilateral stroke and any perioperative stroke or death, the survival curves in the Figure cross near 10 months and become significantly reduced in the surgical group by 3 years (*P* < .05).

Table 5 summarizes the results for secondary end points. Ipsilateral stroke or death by patient or perioperative stroke or death were reduced by 56% (95% CI, 36% to 76%), and the event rate was 3.6% to 6.5% between groups. However, arterial stenosis was not significant (*P* = .1) even with pericardial infarction (3.6%, compared with 5.1%). However, arterial stenosis was reduced by 56% (95% CI, 36% to 76%), compared with 7.9% for men

ue y Risk of Risk oup	P.
72)	.004
3.72)	.12
70)	<.001
3.52)	.09
3.62)	.26
3.37)	.08
3.39)	.16


cal physician included six cerebral in- n of the ran- ebral infarc- distribution, stroke.

significance 2.7 years of s having com- 3; 68%, 2; and ecause surgi- t risk during arterectomy, patients was ars, compari- restated the years. Table ber of events estimates pre- 1 followed for ar risk of ip- perioperative r the medical

rgical group. ateral stroke as 53% of the edical group value for the en the treat- c of primary e-sample test

For the pri- l stroke and r death, the re cross near nificantly re- o by 3 years

y end points although not ant (Table 3 . or stroke or e, or death— oint for the tion in 5-year 35% CI, 39% ke or death,



Proportion of patients without end point at a given time during follow-up, by treatment group, using Kaplan-Meier estimation method. Solid line indicates medical patients; broken line, surgical patients; and TIA, transient ischemic attack.

the surgery group had a 20% reduction in events (95% CI, -2% to 37%). The reduction due to CEA in major ipsilateral stroke or perioperative death was 43% (95% CI, -17% to 72%). Table 4 shows the causes of death were similar for the two groups, except for death not proved to result from myocardial infarction.

Table 5 summarizes results for ipsilateral stroke or any perioperative stroke or death by patient subgroups. In men, CEA reduced the 5-year event rate by 66% (95% CI, 36% to 82%); in women, the event rate was reduced by 17% (95% CI, -96% to 65%). However, the difference between genders was not statistically significant (P=.10). The proportion of women with perioperative complications was 3.6%, compared with 1.7% for men (P=.12). However, among patients who had no perioperative event, 5-year risk was reduced by 56% for women (95% CI, -50% to 87%), compared with a reduction of 79% for men (95% CI, 52% to 91%) (data

not shown). Table 5 shows a larger risk reduction due to CEA for younger patients, but the difference is not statistically significant (P=.50).

Reanalysis excluding the 146 cross-overs, ie, restricted to those patients who received the assigned treatment, shows that surgery reduced 5-year stroke risk by 55% (95% CI, 23% to 74%). Alternatively, for the 1155 patients who had no contralateral TIA, stroke, or endarterectomy prior to randomization, surgery reduced the 5-year stroke risk by 46% (95% CI, 0% to 71%) (Table 5).

The percentage of stenosis for the 642 patients who received an arteriogram within 6 months preceding randomization is shown in Table 5. Throughout the three groups, ie, for patients with 60% to 69%, 70% to 79%, and 80% to 99% stenosis, there was no statistically significant gradation in reduction of 5-year risk of primary event, but sample sizes were small. The patients for this comparison

Table 4.—Number of Deaths, Overall and by Treatment Group, in the ACAS, December 1987 Through December 31, 1994\*

Cause of Death	Treatment Group	
	Surgical	Medical
Perioperative only	3	1
Ipsilateral stroke	2	3
Stroke on contralateral side or posterior circulation	1	2
Cerebral hemorrhage	3	4
Acute myocardial infarction	21	24
Other cardiac disease	15	24
Other vascular disorder	1	2
Respiratory failure	10	9
Cancer	15	13
Tuberculosis	1	0
Renal failure	4	2
Gastrointestinal system disease	4	3
Nervous system disease	1	0
Trauma	2	0
Unknown	0	2†
<b>Total Deaths</b>	<b>83</b>	<b>89</b>
Deaths per 100 person-years of follow-up	3.5	3.8

\*ACAS indicates Asymptomatic Carotid Atherosclerosis Study.

†One patient died at home, the other at a non-ACAS facility. The patient coordinators were unable to obtain information as to the causes of deaths.

all had an arteriogram performed before randomization, so the risk of stroke from undergoing an arteriogram was not included in the calculations. If a 1.2% arteriogram risk were added to the surgery groups at the three stenosis levels, the risk reductions become 0.35, 0.49, and 0.13, respectively, which are consistent with the overall results of the ACAS.

**COMMENT**

The ACAS was designed to test the efficacy of CEA for preventing ipsilateral stroke during a 5-year period. Even though this report includes patients followed up for a median of only 2.7 years, with 9% having completed 5 years of follow-up, the data demonstrate a statistically significant (P=.004) difference between the estimated 5-year ipsilateral stroke rates of 11.0% for the medical and 5.1% for the surgical group. Moreover, the results are in the same direction for all subgroups considered, including deciles of stenosis (although not statistically significant because of small sample size), and for various secondary cerebrovascular end points. Furthermore, the results are virtually the same when restricted to all patients receiving the assigned treatment, and are almost identical for patients without previous contralateral symptoms or endarterectomy.

Approximately 70% of our medical and surgical patients had arteriographic stenoses less than 80%. Even so, the estimated 5-year ipsilateral stroke rate in the ACAS medical group was 11.0% (about 2.3% annually). The stroke rate in the medically managed group decreased to the lower end of the previously reported range, perhaps as a re-



**Action Due  
ary in 5-y Risk  
ortion of Risk  
edical Group**

<b>95% CI</b>
0.22 to 0.72
...
0.36 to 0.82
...
-0.96 to 0.65
...
0.11 to 0.82
...
-0.07 to 0.70
...
0.00 to 0.71
...
0.13 to 0.86
...
0.23 to 0.74
...
-0.70 to 0.82
...
-0.65 to 0.94
...
-2.19 to 0.91
...

10%.  
vent surgery  
as part of the  
olute risk re-  
om 11.0% to  
lity require-  
necessary to  
years.<sup>25</sup> This  
it subsets at  
e identified.  
d with a low  
elderly pa-  
some ACAS  
hout arteri-  
vasive stud-  
ge patients  
<sup>35</sup> These and  
ice costs if

ated that the  
tion can be  
ingent qual-  
lucose surgical  
major reason  
nd mortality  
to have been  
had under-  
of the study.  
omplications  
e improved  
ociated with

contrast arteriography. The ACAS has established that men with a good life expectancy who have asymptomatic carotid artery stenosis with at least 60% reduction in diameter are protected from stroke by CEA, whereas the results for women are less certain. Following CEA, the relative stroke risk reduction for men and women combined is 53%, with an absolute 5-year risk reduction from 11.0% to 5.1%. The 5-year reduction in stroke risk among men was 66% and among women 17%, perhaps because of the higher perioperative complication rate in women. Excluding arteriographic and perioperative complications, the risk reduction was 79% for men and 56% for women.

The Asymptomatic Carotid Atherosclerosis Study clinical trial was supported by an investigator-initiated research grant (RO1 NS22611) from the US Public Health Service National Institute of Neurological Disorders and Stroke.

The authors acknowledge the volunteerism of all patients in the Asymptomatic Carotid Atherosclerosis Study.

**Funding Agency.**—National Institute of Neurological Disorders and Stroke, Bethesda, Md; Michael D. Walker, MD; John R. Marler, MD; Murray Goldstein, DO; Patricia A. Grady, PhD.

**Executive Committee.**—James F. Toole, MD, chair; William H. Baker, MD; John E. Castaldo, MD; Lloyd E. Chambless, PhD; Wesley S. Moore, MD; James T. Robertson, MD; Byron Young, MD; Virginia J. Howard, MSPH (ex officio).

**Executive Committee Liaison.**—John R. Marler, MD.

**Operations Center.**—Bowman Gray School of Medicine of Wake Forest University, Winston-Salem, NC; James F. Toole, MD; Virginia J. Howard, MSPH; Suzanne Purvis; Dee Dee Vernon; Kelley Needham; Pam Beck; Marty Dozier; David S. Lefkowitz, MD; George Howard, DrPH; John R. Foubert III, MD; David M. Herrington, MD; Curt D. Furberg, MD, PhD; Karla Essick; Ralph M. Hicks, Jr, MA, MEd.

**Statistical Coordinating Center.**—University of North Carolina at Chapel Hill; Lloyd E. Chambless, PhD; J. J. Nelson, MSPH; Will Ball; Ernestine Bland; Sean Condon, MS; Thomas Elliott, MS (deceased); James E. Grizzle, PhD; Dick Hayes, MCJ; Scott Henley; Jeffrey Johnson, MSPH; James Locklear, MS; Margaret S. Misch, MS; Catherine C. Paton, MSPH; Skai Schwartz, MA; Climmon Walker; O. Dale Williams, PhD.

**Data and Safety Monitoring Committee.**—J. Donald Easton, MD, chair; Jerry Goldstone, MD; John M. Hallenbeck, MD; Julian T. Hoff, MD; Herbert R. Karp, MD; Richard A. Kronmal, PhD.

**Cranial Computerized Tomography Reading Center.**—University of Cincinnati (Ohio); Thomas G. Brott, MD; Thomas A. Tomsick, MD; Joseph Broderick, MD; Laura Sauerbeck, RN; Christine Blum, RN.

**Cerebrovascular End Point Review Committee.**—Mark Dyken, MD, chair; John C. M. Brust, MD; Arthur R. Dick, MD; Robert A. Gotshall, MD; Albert Heyman, MD; Phillip D. Swanson, MD.

**Publications Committee.**—James F. Toole, MD, chair; Harold P. Adams, Jr, MD; Lloyd E. Chambless, PhD; Robert Dempsey, MD; Calvin B. Ernst, MD; John Rothrock, MD (with special appreciation to Stanley N. Cohen, MD, for desk editing).

**Participating Centers (in order of number of patients randomized, number of patients in brackets, and personnel listed in the following order: Principal Investigator, Co-Investigator, Coordinator, and Other Key Personnel, Current and Past)**

**Lehigh Valley Hospital,** Allentown, Pa [142]; John Castaldo, MD; Gary Nicholas, MD; Joan Longenecker; Peter Barbour, MD; Alan Berger,

MD; Victor J. Celani; Nancy Eckert, RN; James Goodreau, MD; Judith Hutchinson, RN; Donna Jenny, RN; Zwu-Shin Lin, MD; Alice Madden, RN; James L. McCullough, MD; Kenneth McDonald, MD; William Pistone, MD; Alexander D. Rae-Grant, MD; James Redenbaugh, MD; James Rex, MD; Christopher J. Wohiberg, MD.

**Marshfield Clinic, Wisconsin** [121]; Percy Karanjia, MD; Mark Swanson, MD; Sandra Lobner, LPN; R. Lee Koits, MD; Marvin E. Kuehner, MD; Bradley C. Hiner, MD; Kenneth P. Madden, MD; Robert D. Carlson, MD; J. Steven Davis, MD; Thomas Gallant, MD; John J. Warner, MD; Ann Faust, RVT; Nancy Fryza, RVT; John Hasenauer, RVT; Mary Regner, RVT; Linette Ronkin, RVT; Sharon Schaefer, RVT; Dawn Strack, RVT; Lynn C. Turner, RVT; Ann Walgenbach, RVT; Julie Graves; Sara Michalski; Linda Schuette.

**Columbia University,** New York, NY [91]; J. P. Mohr, MD; Donald Quest, MD; Annette Cruz; Ralph Sacco, MD; T. K. Tatemichi, MD; Randolph Marshall, MD; Henning Mast, MD; Oscar Ramos, MD; James Correll, MD; Richard Libman, MD; George Petty, MD; Anilda Cabrere, MD; Lorraine Oropeza, RN, RVT; Tina Gonzalez.

**University of Kentucky Chandler Medical Center,** Lexington [88]; Byron Young, MD; Creed Pettigrew, MD; Ruthie Sadler, RN; Robert Dempsey, MD; Eric Edeand, MD; Jerry Sherrow, RVT; Marcie Hauer, RN; Charles Lee, MD; Jane Norton; Michael McQuillen, MD; Sally Mattingly, MD; Steven Dekosky, MD; Andrew Massey, MD.

**Hôpital de l'Enfant Jésus,** Québec City, Québec [86]; Denise Simard, MD; Jean-François Turcotte, MD; Charlotte Benguigui, RN; Jacques Côté, MD (deceased); Jean-Marie Bouchard, MD; Claude Roberge, MD; Denis Brunet, MD; Fernand Bédard; Réginald Langelier, MD; Marcel Lajeunesse; Jean Marc Bigaquette; Jasmin Parent.

**University of California at San Diego** [78]; Patrick Lyden, MD; Robert Hye, MD; Stacy Lewis, RN; Gerry Cali, RN; Traci Babcock, RN; Barbara Taft-Alvarez, RN; John Rothrock, MD; Mark Brody, MD; Richard Zweifler, MD; Mark Sedwitz, MD; Bruce Stabile, MD; Julie A. Freischlag, MD; Yehuda G Wolf, MD; James Sivo; John Forsythe; Melody Adame.

**Loyola University Medical Center,** Maywood, Ill [77]; William H. Baker, MD; Sudha Gupta, MD; Katy Burke; Howard P. Greisler, MD; Fred N. Littooy, MD; Michael A. Kelly, MD.

**University of Tennessee,** Memphis [74]; James T. Robertson, MD; William Pulsinelli, MD; Judith A. Campbell, RN, BSN; John Crockarell, MD; Clarence Watridge, MD; J. Acker, MD; Samard Erkulwater, MD; Michael Jacewicz, MD; Gail Walker; Patrick O'Sullivan, MD; Curtis Sauer, MD; Ken Vasu, MD; Kenneth Gaines, MD; Bijan Bakhtian, MD; Tulio E. Bertorini, MD; Susan Bennett, PA-C; Terrye Thomas, BSN; Nan Stahl, RN; Connie Taylor, RN; Mary Ann Giampapa; Joan Connell; Judy Riley; Angeleah Bradley; Jeanette Davis, RN; Kathy Newman; Rebecca Manning; Marcella McCrea.

**University Hospital,** London, Ontario [63]; Vladimir Hachinski, MD; Gary Ferguson, MD; Cheryl Mayer, RN; Henry J. M. Barnett, MD; S. J. Peerless, MD; A. M. Buchan, MD; Howard Reichman, MD; Andrew Kertesz, MD; Stephen Lownie, MD; Carole White, RN; Allan Fox, MD; Richard Rankin, MD.

**Victoria Hospital,** London, Ontario [61]; J. David Spence, MD; H. W. K. Barr, MD; Leslie Paddock-Eliasziw, RN; L. J. P. Assis, MD; J. H. W. Pexman, MD; Maria DiCicco, RVT; Barb Tate, RN; Caroline James, RVT.

**Virginia Mason Clinic,** Seattle, Wash [57]; Edmond Raker, MD; James Coatsworth, MD; Sandy Harris, RN; Hugh Beebe, MD; Richard Birchfield, MD; Kathy Butler-Levy; Robert Crane, MD; David Fryer, MD; James MacLean, MD; Laird Patterson, MD; Terence Quigley, MD; John Ravits, MD; Lynne Taylor, MD; Stacy Pullen, RDMS; Shannon Boswell, RDMS; Karen Kenny, RDMS, RVT.

**University of Cincinnati Ohio** [55]; Thomas Brott, MD; Joseph Broderick, MD; Thomas Tomsick, MD;

Laura Sauerbeck, RN; L. F. Roedersheimer, MD; Richard Fowl, MD; John Tew, MD; Richard Kempczinski, MD; Robert Reed, MD; Richard Welling, MD; Christine Blum, RN; Bill Schomaker, RN.

**Bowman Gray School of Medicine of Wake Forest University,** Winston-Salem, NC [52]; David Lefkowitz, MD; J. Michael McWhorter, MD; Charles L. Branch, MD; Jean Satterfield; Robert Cordell, MD; Richard Dean, MD; George Plonk, MD; Gary Harpold, MD; James F. Toole, MD; Francis Walker, MD; Cathy Nunn, RN RVT; Larry Myers, MBA, RDMS, RVT; Charles Tegeler, MD; Sharon Hardin, RVT, RCMS; Dana Meeds, RT-R, RVT.

**University of Iowa Hospital and Clinics,** Iowa City [49]; Harold P. Adams, Jr, MD; Christopher M. Loftus, MD; Lynn A. Vining, RN; Birgette H. Bendixen, MD, PhD; José Biller, MD; John D. Corson, MD; Patricia H. Davis, MD; John C. Godersky, MD; David L. Gordon, MD; Michael R. Jacoby, MD; Laurens J. Kappelle, MD; Timothy F. Kresowik, MD; E. Eugene Marsh III, MD; Betsy B. Love, MD; Asad R. Shamma, MD; Karla J. Grimsmann, RN; Dawn M. Karboski, RN; Ed V. Miller, RVT.

**Johns Hopkins Bayview Medical Center,** Baltimore, Md [46]; Constance Johnson, MD; Calvin Jones, MD; Brenda Stone, CRNP; Mairi Pat Maguire, RN; Christopher Earley, MD; Peter Kaplan, MD; John Cavaluzzi, MD; Gerald Waters, PA-C; Betty Chachich, RN.

**St John's Mercy Medical Center,** St Louis, Mo [44]; Arthur Auer, MD; William Logan, MD; Mary Wilcox, RN, BSN; Barbara Green, MD; Joseph Hurley, MD; Richard Pennell, MD; John Woods, MD; Richard Levine, MD; James Nepute, MD; Jeffrey Thomasson, MD; Claire Blackburn, RN, RVT; Marcia Foldes, BSN, RVT; Kathy Klemp, RN, RVT; Becky Nappier, RT, RVT; Karen Rutherford, BSN, RVT; Sally Schroer, RN, BSN; Jena Hogan, RN, RVT; Lynne Thorpe, RN, RVT.

**University of Arizona Health Sciences Center,** Tucson [42]; William M. Feinberg, MD; Glenn C. Hunter, MD; Denise C. Bruck, AAS; Victor M. Bernhard, MD; Kenneth E. McIntyre, MD; L. Philip Carter, MD; Scott S. Berman, MD; Joseph L. R. Mills, MD; Enrique L. Labadie, MD; Darry S. Johnson, MD; Constantine G. Moschonas, MD; Robert H. Hamilton, MD; Scott C. Forrer, MD; Joachim F. Seeger, MD; Raymond F. Carmody, MD; Brenda K. Vold, RN, BSN; Richard L. Carlson; Joan E. Laguna; John P. Krikawa; Jenifer J. Devine, RN, BSN, RVT; Amalia M. Castrillo; Sheryl L. Kistler, RN, RVT; Betty Ledbetter; Kathleen (Sue) Dorr.

**University of Mississippi Medical Center,** Jackson [39]; Robert R. Smith, MD; Armin F. Haerer, MD; Robin L. Brown, RN, BSN; William Russell, MD; Edward Rigdon, MD; Robert Rhodes, MD; Evelyn Smith, RVT; Michael Graeber, MD; David Doorenbos, MD; S. H. Subramony, MD; David L. Gordon, MD.

**Milton S. Hershey Medical Center,** Hershey, Pa [35]; Robert G. Atnip, MD; Robert W. Brennan, MD; Diane Friedman, RN, MSN; Marsha M. Neumyer, RVT; Brian L. Thiele, MD; Florence Smith, RN; John D. Barr, MD; R. Bradford Duckrow, MD; Cindy Janesky, MD; Jon W. Meilstrup, MD; Kevin P. McNamara, MD; Lawrence D. Rodichok, MD; Leslie Stewart, MD; Maureen Sullivan, MD; Mark Wengrovitz, MD.

**University of Texas Southwestern Medical Center,** Dallas [35]; G. Patrick Clagett, MD; Hal Unwin, MD; Wilson Bryan, MD; Chris Matkins; Carolyn Patterson; Candy Alway, RDMS; Patty Boyd; Mary Inman, RN; Christie Albiston; Eva Scoggins; John Swilling.

**University of California at Los Angeles** [35]; Wesley S. Moore, MD; Stanley N. Cohen, MD; Kathleen G. Walden, RN; Samuel S. Ahn, MD; Edwin C. Amos, MD; J. Dennis Baker, MD; Bruce H. Dobkin, MD; Carlos E. Donayre, MD; Julie A. Freischlag, MD; Hugh A. Gelabert, MD; Sheldon E. Jordan, MD; Herbert I. Machleder, MD; William J. Quinones-Baldrich, MD; Jeffrey L. Saver, MD; Suzie M. El-Saden, MD; Richard C. Holgate, MD; Bradley A. Jabour, MD; J. Bruce Jacobs, MD; Theresa M. Abraham, RN; Candace L. Vescera,

RN; Jeanine A. von Rajcs, RN; Vicki L. Carter, RN, RVT; Dale T. Carter, RVT; De Ette Dix-Goss, RVT; Eugene C. Hernandez, RVT.

*Oregon Health Sciences University*, Portland [34]: Lloyd Taylor, MD; Bruce Coull, MD; Letha Lobo, RN, RVT; Gregory Moneta, MD; John Porter, MD; Richard Yeager, MD; Lucy Whittaker, RN.

*Yale University*, New Haven, Conn [34]: Lawrence M. Brass, MD; Richard J. Gusberg, MD; Anne M. Lovejoy, PA-C; Pierre B. Fayad, MD; Bauer Sumpio, MD, PhD; George H. Meier, MD; Vicky M. Chang, ANP, MSN; Karen Marzittelli, ANP, MSN; Douglas Chyatte, MD; Lynwood Hammers, DO; Fran Lepore, RDMS, RT; Frank J. Pavalkis, PA-C; Jill Mele; Donna Kisiel, PA-C.

*University of Arkansas for Medical Sciences*, Little Rock [32]: Robert W. Barnes, MD; Michael Z. Chesser, MD; R. Lee Archer, MD; Bernard W. Thompson, MD; Colette MacDonald; Gary W. Barone, MD; John F. Eldt, MD; David Harshfield, MD; David McFarland, MD; Jess R. Nickols, MD; E. Carol Howard, BSN, RN, RVT; M. Lee Nix, BSN, RN, RVT; Judy Kaye Overstreet, RN, RVT; Rhonda Troillet, RDMS, RVT.

*Medical College of Virginia (Virginia Commonwealth University)*, Richmond [27]: John Taylor, MD; H. M. Lee, MD; Patricia Akins; John W. Harbison, MD; Rhonda M. Pridgeon, MD; Warren L. Felton, MD; Marc Posner, MD; M. Sobel, MD; Guy Clifton, MD; Chris Conway; Anne Cockrell; Warren Stringer, MD; Jean Wingo; Brenda Nichols; Wendy Smoker, MD; Ruth Fisher.

*Barrow Neurological Institute at St Joseph's Hospital and Medical Center*, Phoenix, Ariz [26]: Robert F. Spetzler, MD; James L. Frey, MD; Joseph M. Zabramski, MD; Sonna Lea Hunsley, RN; Heidi Jahnke, BSN; Kathryn L. Plenge, MD; Rosemary Holland, RVT; Roberta Turner, RVT; Duane Strava, RVT; Susan Stumpff, RVT; John Hodak, MD; Richard A. Flom, MD; Bruce L. Dean, MD; Richard A. Thompson, MD.

*Ochsner Clinic*, New Orleans, La [21]: Richard Hughes, MD; Bruce Lepler, MD; John Bowen, MD; Cheryl Benoit, CCRC; Larry Hollier, MD; John Ochsner, MD; Richard Strub, MD; Vickie Lang; Vicki Cahanin.

*University of Medicine and Dentistry of New Jersey*, Newark [20]: Robert W. Hobson II, MD; Fred Weisbrod, MD; Stephen Kamin, MD; Tom Back; Zafar Jamil, MD; Carolyn Rogers; Bea Lains; Larry Hart.

*New England Medical Center*, Boston, Mass [19]: Louis Caplan, MD; Thomas O'Donnell, MD; Loretta Barron, RN; Michael Pessin, MD; Dana DeWitt, MD; William Mackey, MD; Michael Belkin, MD; Robert McLaughlin, RVT; Paula Heggerick, RVT.

*Henry Ford Hospital*, Detroit, Mich [17]: Calvin B. Ernst, MD; K. M. A. Welch, MD; Judith Wilczewski, RN; Wendy Robertson, PA-C; Sheila Daley, RN; Joseph P. Elliot, Jr, MD; Daniel J. Reddy, MD; Alexander D. Shephard, MD; Roger F. Smith, MD; Steven Levine, MD; Nabih Ramadan, MD; Gretchen Tietjen, MD; Panayiot Mitsias, MD; Mark Gorman, MD; Michalene McPharlin, RN; Suresh Patel, MD; Rajeev Deveshwar, MD; Nora Lee, MD; James Kokkinos, MD.

*University of New Mexico*, Albuquerque [13]: Asklie Bruno, MD; Eric Weinstein, MD; James Kunkel, MD (deceased); Anna Kratochvil, RN; Edie Johnson, RN, MSN, FNP; Susan Steel, RN.

*Sunnybrook Health Science Centre*, University of Toronto, North York, Ontario [11]: John Norris, MD; David Rowed, MD; Beverley Bowyer, RN; Marek Gawel, MD; Perry W. Cooper, MD; Dianne Brodie, RVT.

*Harbin Clinic*, Rome, Ga [10]: John S. Kirkland, MD, PhD; Jay A. Schecter, MD; Nell W. Farrar, PA-C; Raymond Capps, MD; E. Leon Rhodes, MD; D. Michael Rogers, MD; Jeffrey T. Glass, MD; Robert Naguszewski, MD; William Naguszewski, MD; Bobbie Maddox; Bobbie Dollison; Lynda Moulton; Polly Cole; Paul Kinsella; Alisa Ansley; Noel Britz.

*Roanoke Neurological Associates*, Virginia [10]: Don H. Bivins, MD; Edwin L. Williams II, MD; Jesse T. Davidson III, MD; William Elias, MD;

Donna Atkins, RN; Phyllis B. Turner, RN, RVT; J. Gordon Burch, MD; Donald B. Nolan, MD; Renee Speese; Candace D. Foley, RN.

*Singing River Hospital*, Pascagoula, Miss [6]: Terrence J. Millette, MD; Dewey H. Lane, MD; Cynthia Almond, RN, RVT; Roland J. Mestayer III, MD.

*California Pacific Medical Center*, San Francisco [4]: Phillip Calanchini, MD; Robert Szarnicki, MD; Pat Radosevich; Linda Elias; Peggy McCormick; Charles Gould, MD; Forbes Norris, MD; Eric Denys, MD; Robert Bernstein, MD; Donna Dubono; Keith Atkinson; Mark Peters.

*Northwestern University Medical School*, Chicago, Ill [4]: Bruce Cohen, MD; James Yao, MD; Susan Roston, RN; Donna Blackburn, RN; José Biller, MD; Jeffrey Saver, MD; Linda Chadwick, RN; Douglas Chyatte, MD; Walter McCarthy, MD; William Pearce, MD; Jeff Frank, MD; Ernesto Fernandez-Beer, MD; James Patrick, MD.

*University of Rochester*, New York [3]: Richard Green, MD; Richard Satran, MD; John Ricotta, MD; James DeWeese, MD; Joshua Hollander, MD; Mollie O'Brien, RN; JoAnne McNamara, RN; Sandra Rose, RT, RDMS; Dahne Cohen, MD.

*Cleveland Clinic*, Ohio [1]: Anthony Furlan, MD; John Little, MD; Bernadine Bryerton, RN; Cathy Sila, MD; Isam Awad, MD; Marc Chimowitz, MD; Sylvia Robertson, ARRT, RDMS; Cindy Becker, ARRT, RDMS; David Paushter, MD.

*Arteriogram Quality Control*.—Bruce L. Dean, MD; Daniel H. O'Leary, MD.

*Doppler and OPG-Gee Quality Control*.—Anne M. Jones, BSN, RVT, RDMS; John J. Ricotta, MD; William Gee, MD; Nancy Shebel, BSN, RN, RVT.

*Pathology*.—Mark J. Fisher, MD; Eric A. Schenk, MD.

*TIA/Stroke Validation*.—Anthony J. Furlan, MD; Nancy N. Futrell, MD; Michael Kelly, MD; Clark H. Millikan, MD.

*General Consultants*.—H. C. Diener, MD; William S. Fields, MD; Marshall F. Folstein, MD; Jean-Claude Gautier, MD; Michael J. G. Harrison, MD; William K. Hass, MD; Michael G. Hennerici, MD; Richard Satran, MD; Merrill P. Spencer, MD; Gerhard M. von-Reutern, MD.

## References

1. Timsit SG, Sacco RL, Mohr JP, et al. Early clinical differentiation of cerebral infarction from severe atherosclerotic stenosis and cardioembolism. *Stroke*. 1992;23:486-491.
2. Hennerici M, Hülsbømer H-B, Hefter H, Lammerms D, Rautenberg W. Natural history of asymptomatic extracranial arterial disease. *Brain*. 1987; 110:777-791.
3. Autret A, Saudeau D, Bertrand PH, Pourcelot L, Marchal C, DeBoisvilliers S. Stroke risk in patients with carotid stenosis. *Lancet*. 1987;1:888-890.
4. O'Hollerhan LW, Kennelly MM, McClurken M, Johnson JM. Natural history of asymptomatic carotid plaque. *Am J Surg*. 1987;154:659-662.
5. Norris JW, Zhu CZ, Bornstein NM, Chambers BR. Vascular risks of asymptomatic carotid stenosis. *Stroke*. 1991;22:1485-1490.
6. Jorgensen L, Torvik A. Ischaemic cerebrovascular diseases in an autopsy series, 2. *J Neurol Sci*. 1969;9:285-320.
7. Cote R, Barnett HJM, Taylor DW. Internal carotid occlusion. *Stroke*. 1983;14:898-902.
8. Nicholls SC, Kohler TR, Bergelin RO, Primozieh JP, Lawrence RL, Strandness DE Jr. Carotid artery occlusion. *J Vasc Surg*. 1986;4:479-485.
9. Moore WS, Barnett HJM, Beebe HG, et al. Guidelines for carotid endarterectomy. *Stroke*. 1995;26:188-201.
10. Riles TS, Adelman MA, Gibstein LA. Incidence of premonitory transient ischemic attacks before ischemic strokes. In: Veith FJ, ed. *Current Critical Problems in Vascular Surgery*. St Louis, Mo: Quality Medical Publishing Inc; 1994;6:264-268.
11. Till JS, Toole JF, Howard VJ, Ford CS, Williams D. Declining morbidity and mortality of carotid endarterectomy. *Stroke*. 1987;18:823-829.
12. Wilson SE, Mayberg MR, Yatsu FM. Defining

the indications for carotid endarterectomy. *Surgery*. 1988;104:932-933.

13. The Asymptomatic Carotid Atherosclerosis Study Group. Study design for randomized prospective trial of carotid endarterectomy for asymptomatic atherosclerosis. *Stroke*. 1989;20:844-849.

14. Howard G, Chambless LE, Baker WH, et al. A multicenter validation study of Doppler ultrasound versus angiogram. *J Stroke Cerebrovasc Dis*. 1991; 1:166-173.

15. Brott T, Tomsick T, Feinberg W, et al, for the Asymptomatic Carotid Atherosclerosis Study Investigators. Baseline silent cerebral infarction in the Asymptomatic Carotid Atherosclerosis Study. *Stroke*. 1994;25:1122-1129.

16. Moore WS, Vecsera CL, Robertson JT, Baker WH, Howard VJ, Toole JF. Selection process for surgeons in the Asymptomatic Carotid Atherosclerosis Study. *Stroke*. 1991;22:1353-1357.

17. Folstein MF, Folstein SE, McHugh PR. 'Minimal State': a practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res*. 1975;12:189-198.

18. Lefkowitz DS, Brust JC, Goldman L, et al. A pilot study of the end point verification system in the Asymptomatic Carotid Atherosclerosis Study. *J Stroke Cerebrovasc Dis*. 1992;2:92-99.

19. Jennett B, Bond M. Assessment of outcome after severe brain damage: a practical scale. *Lancet*. 1975;1:480-484.

20. Hobson RW II, Weiss DG, Fields WS, et al, for the Veterans Affairs Cooperative Study Group. Efficacy of carotid endarterectomy for asymptomatic carotid stenosis. *N Engl J Med*. 1993;328:221-227.

21. North American Symptomatic Carotid Endarterectomy Trial Collaborators. Beneficial effect of carotid endarterectomy in symptomatic patients with high-grade stenosis. *N Engl J Med*. 1991;325:445-453.

22. Kalbfleisch JD, Prentice RL. *The Statistical Analysis of Failure Time Data*. New York, NY: John Wiley & Sons Inc; 1980:14.

23. Lachin JM. Statistical properties of randomization in clinical trials. *Control Clin Trials*. 1988; 9:289-311.

24. O'Brien PC, Fleming TR. A multiple testing procedure for clinical trials. *Biometrics*. 1979;35:549-556.

25. Laupacis A, Naylor CD, Sackett DL. How should the results of clinical trials be presented to clinicians? *ACP J Club*. May/June 1992:A12-A14.

26. CASANOVA Study Group. Carotid surgery versus medical therapy in asymptomatic carotid stenosis. *Stroke*. 1991;22:1229-1235.

27. Mayo Asymptomatic Carotid Endarterectomy Study Group. Results of a randomized controlled trial of carotid endarterectomy for asymptomatic carotid stenosis. *Mayo Clin Proc*. 1992;67:513-518.

28. Halliday AW, Thomas D, Mansfield A. The Asymptomatic Carotid Surgery Trial (ACST): rationale and design. *Eur J Vasc Surg*. 1994;8:703-710.

29. European Carotid Surgery Trialists' Collaborative Group. European Carotid Surgery Trial: interim results for symptomatic patients with severe (70-99%) or with mild (0-29%) carotid stenosis. *Lancet*. 1991;337:1235-1243.

30. Toole JF, Castaldo JE. Accurate measurement of carotid stenosis: chaos in methodology. *J Neuroimaging*. 1994;4:222-230.

31. Rothwell PM, Gibson RJ, Slattery J, Sellar RJ, Warlow CP, for the European Carotid Surgery Trialists' Collaborative Group. Equivalence of measurements of carotid stenosis. *Stroke*. 1994;25:2435-2439.

32. Alexandrov AV, Bladin CF, Magissano R, Norris JW. Measuring carotid stenosis: time for a reappraisal. *Stroke*. 1993;24:1292-1296.

33. Horn M, Michelini M, Greisler HP, Littooy FN, Baker WH. Carotid endarterectomy without arteriography. *Ann Vasc Surg*. 1994;8:221-224.

34. Chervu A, Moore WS. Carotid endarterectomy without arteriography. *Ann Vasc Surg*. 1994;8:296-302.

35. Morasch MD, Hodgett D, Burke K, Baker WH. Selective use of the intensive care unit following carotid endarterectomy. *Ann Vasc Surg*. In press.

# Efficacy of Diet in Cholesterol Lipoproteins: The Diet Writing Group

**Objective.**—fat, saturated (LDL-C) levels  
**Design.**—S Participant with LDL-C levels for age a usual care group  
**Intervention.** 28% of energy unsaturated fat to exceed 15C  
**Main Outcome.** LDL-C level at 3 year and mean included red b high-density lip triglyceride lev  
**Results.**—/creased signif (all  $P < .001$ ). L by 0.40 mmol/l ing for baseline ference between interval [CI], — ( $P = .02$ ). There height or seru  
**Conclusion.** levels over 3 y equacy, and p cence.

On behalf of the Group for this article. Eva Obarzanek, Sue Y. S. Kimm, M. A. Barton, PhD, Linc. Frank A. Franklin, I. Morton, MD, PhD. Reprint requests (and Medical Rese. Baltimore, MD 212